

Topic Area: Occupational Safety and Health

OSH-1: Reduce the mortality rate due to work-related injuries (unintentional and homicide)

OSH-1a: All industries

Target:	1.7 per 100,000 employed persons
Baseline: (Year)	2.1 per 100,000 employed persons (2008)
Data source:	New Jersey Census of Fatal Occupation Injuries, Consumer, Environmental, and Occupational Health Service, New Jersey Department of Health

OSH-1b: Construction

Target:	8.8 per 100,000 construction workers
Baseline: (Year)	11.0 per 100,000 construction workers (2008)
Data source:	New Jersey Census of Fatal Occupation Injuries, Consumer, Environmental, and Occupational Health Service, New Jersey Department of Health

OSH-2: Reduce deaths due to pneumoconiosis among persons 15 years and older

Target:	6.9 per 1,000,000 standard population aged 15 years and older (age-adjusted)
Baseline: (Year)	6.0 per 1,000,000 standard population aged 15 years and older (age-adjusted) (2007)
Data source:	Consumer, Environmental, and Occupational Health Service, New Jersey Department of Health

OSH-3: Reduce elevated (≥ 25 $\mu\text{g/dL}$) blood lead concentration from work exposures among adults age 17 years and older

Target:	1.6 per 100,000 employed persons
Baseline: (Year)	1.8 per 100,000 employed persons (2007)
Data source:	Adult Blood Lead Epidemiology and Surveillance, Consumer, Environmental, and Occupational Health Service, New Jersey Department of Health